


RECENT RESULTS FROM KTeV

L. BELLANTONI
for the KTeV Collaboration
*Fermi National Accelerator Laboratory,
Batavia IL 60510, U.S.A.*



The implications of the published KTeV $K_L^0 \rightarrow \pi^0 \mu^+ \mu^-$ result for interpreting recent $\Sigma^+ \rightarrow p \mu^+ \mu^-$ results are discussed. The status of the KTeV $\Xi^0 \rightarrow \Sigma^+ \mu^- \nu$ analysis is given. The KTeV $|V_{us}|$ result is also given.

1 $K_L^0 \rightarrow \pi^0 \mu^+ \mu^-$

The decay $K_L^0 \rightarrow \pi^0 \mu^+ \mu^-$ has three contributions to the amplitudes: a CP conserving term from intermediate states with two photons; a CP violating term from indirect CP violation; and a direct CP violating term from 2nd-order electroweak penguin and box diagrams. The process continues to be of theoretical interest; see in particular the presentation by Christopher Smith at this conference. There is also interest on the experimental side as a consequence of a recent unusual result from the HyperCP collaboration¹.

In searching for $\Sigma^+ \rightarrow p \mu^+ \mu^-$, HyperCP found three events. This is not so unusual; what is peculiar is that to within the $\sim 0.5 \text{ MeV}$ resolution of the mass measurements, all three events have the same dimuon mass of 214.3 MeV . In the standard model, the muon pair is produced by an off-shell photon and a spread of dimuon masses is expected. The HyperCP collaboration estimates that the probability of three events from an intermediate photon having the same mass to this level of precision is about 0.8%, and they suggest that there may be a new intermediate neutral state causing this anomaly.

This postulated new state would be a flavor changing neutral current when coupling to quarks, but not when coupling to leptons. The scale of the partial width for the three observed events, $\Gamma(\Sigma^+ \rightarrow p P^0, P^0 \rightarrow \mu^+ \mu^-) \sim 2.6 \times 10^{-19} \text{ MeV}$ is too small for a strong interaction, which have widths on the order of a few 10^{-12} MeV in Σ^+ decays. Also of course, the active state of research into new bound QCD states notwithstanding, a new narrow hadronic resonance

in this mass range would be a surprise. So we consider new point-like particles. Allowing that the new interaction conserves parity and angular momentum, the only possible $(J)^P$ values for decays into $\mu^+\mu^-$ are $(0)^-$ and $(1)^-$. However if this P^0 is a vector current then it will also contribute to $K_L^0 \rightarrow \pi^0\mu^+\mu^-$ decay.

The existing KTeV limit² on $\text{Br}(K_L^0 \rightarrow \pi^0\mu^+\mu^-)$ of 3.8×10^{-10} at the 90% C.L., while an order of magnitude above the standard model prediction³ of $(1.5 \pm 0.3) \times 10^{-11}$, does correspond to a partial width of $\Gamma(K_L^0 \rightarrow \pi^0 P^0, P^0 \rightarrow \mu^+\mu^-) \sim 4.8 \times 10^{-24} \text{ MeV}$, nearly 5 orders of magnitude less than the postulated HyperCP rate. The vector current hypothesis is thus disfavored.

Work on a new limit for $\text{Br}(K_L^0 \rightarrow \pi^0\mu^+\mu^-)$ based on the full KTeV data set is in progress.

2 $\Xi^0 \rightarrow \Sigma^+\mu^-\nu$

The first observation⁴ of the decay $\Xi^0 \rightarrow \Sigma^+\mu^-\nu$ was made on the 1997 KTeV dataset; here we report results based on the 1999 data, which corresponds to about 3×10^8 Ξ^0 decays. We reconstruct the Σ^+ hyperon in the $p\pi^0; \pi^0 \rightarrow \gamma\gamma$ mode and normalize the data sample with $\Xi^0 \rightarrow \Lambda\pi^0; \Lambda \rightarrow p\pi^-$. Backgrounds are from Ξ^0 , Λ and $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ decays; for all decays other than $\Xi^0 \rightarrow \Lambda\pi^0; \Lambda \rightarrow p\pi^-$ and $K_L^0 \rightarrow \pi^+\pi^-\pi^0$, Monte Carlo samples corresponding to 10 or more times the data sample have been generated. The kaon background is studied with data events where the highest-momentum track is negatively charged. For $K_L^0 \rightarrow \pi^+\pi^-\pi^0$, high momentum π^+ and π^- are equally probable, but the hyperon signal is overwhelmingly comprised of events with high-momentum positively charged tracks. Background from $\Xi^0 \rightarrow \Lambda\pi^0; \Lambda \rightarrow p\pi^-$ is suppressed by relying on the neutrino in the signal mode to produce a missing momentum component perpendicular to the line of Ξ^0 flight. The overall background level is very low, and there are nine events in the data, as shown in Figure 1. We obtain, as a preliminary result, $\text{Br}(\Xi^0 \rightarrow \Sigma^+\mu^-\nu) = (4.3 \pm 1.4) \times 10^{-6}$.

3 $|V_{us}|$

A long standing issue in flavor physics has been a discrepancy, at the 2σ level, of the measured values of $|V_{ud}|$, $|V_{us}|$ and $|V_{ub}|$ from the unitarity constraint for the first row of the CKM matrix. It should be noted that the first row of the matrix is the one that provides the most stringent test of 3-generation unitarity. The value of $|V_{ud}|$ is precisely known from nuclear and neutron beta decays, and $|V_{ub}|$ is small enough to be irrelevant here. $|V_{us}|$ may be determined from semileptonic kaon decays; in the past only the $K_L^0 \rightarrow \pi^0 e^\pm \nu$ mode has been used, due to uncertainty in the form factors of the $K_L^0 \rightarrow \pi^0 \mu^\pm \nu$ mode.

To address this discrepancy, KTeV has sought improved measurements of $\Gamma(K_L^0 \rightarrow \pi^0 e^\pm \nu)$ and $\Gamma(K_L^0 \rightarrow \pi^0 \mu^\pm \nu)$. There being no other decay modes that are known to sufficient precision to normalize the data set, we have measured five ratios of partial widths for the modes $K_L^0 \rightarrow \pi^0 e^\pm \nu$, $K_L^0 \rightarrow \pi^0 \mu^\pm \nu$, $K_L^0 \rightarrow \pi^+\pi^-\pi^0$, $K_L^0 \rightarrow 3\pi^0$, $K_L^0 \rightarrow \pi^+\pi^-$ and $K_L^0 \rightarrow 2\pi^0$. These modes account for nearly all K_L^0 decays, and this fact may be used to extract the branching ratios for the semileptonic modes. With the previously accepted¹⁰ value (see the presentation of Gaia Lanfranchi at this conference) for the lifetime of the K_L^0 , partial widths for the semileptonic modes and then values for $|V_{us}|$ have been extracted.

A full exposition of this work and the related analyses spans a number of publications:

- The radiative corrections - reference⁵
- Measurement of form factors for $K_L^0 \rightarrow \pi^0 e^\pm \nu$ and $K_L^0 \rightarrow \pi^0 \mu^\pm \nu$ - reference⁶
- Check with radiative $K_L^0 \rightarrow \pi^0 e^\pm \nu$ and $K_L^0 \rightarrow \pi^0 \mu^\pm \nu$ decays - reference⁷




Figure 1: Reconstructed $P_{\perp}$ vs. $M_{p\pi^0}$ for $\Xi^0 \rightarrow \Sigma^+ \mu^- \nu$ candidates in the 1999 KTeV data.

- The measurement of the partial width ratios - reference⁸
- The extraction of $|V_{us}|$ from these - reference⁹

The final result, using $f_+(0) = 0.961 \pm 0.008$, is $|V_{us}| = 0.2252 \pm 0.0008_{\text{KTeV}} \pm 0.0021_{\text{ext}}$, where the KTeV uncertainty includes uncertainties in the KTeV branching fractions and form factor measurements, and the ext uncertainty includes uncertainties in $f_+(0)$, K_L^0 lifetime, and radiative corrections. This result does resolve the unitarity discrepancy, and the attendant results show a high degree of internal consistency. While our branching ratios are not in good agreement with the values listed in the 2002 PDG, many of the discrepancies may be explained by postulating that $\text{Br}(K_L^0 \rightarrow \pi^0 e^\pm \nu)$ in the 2002 PDG was too low. In discussion I emphasized that, as noted by Cirigliano *et.al.*¹¹, the PDG did not actually have a precise and direct measurement of $\text{Br}(K_L^0 \rightarrow \pi^0 e^\pm \nu)$ on hand. The value that they recommended was inferred from reports of other measurements and the indisputable constraint that the sum of the branching ratios must be one. It was, if you would, a "global fit", albeit a substantially simpler analysis than many calculations going by that name. Caveat!

4 Acknowledgements

I very much want to thank the organizers of this most excellent conference, and in particular the gracious and highly competent Elizabeth Hautefeuille as well as the sagacious and justly renowned Jean Tran Thanh Van.

The KTeV collaboration gratefully acknowledges the support and effort of the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported in part by the U.S. Dept. of Energy, the National Science Foundation, and the Ministry of Education and Science of Japan.

References

1. H.K. Park *et al*, *Phys. Rev. Lett.* **94**, 021801 (2005).
2. A. Alavi-Harati *et al*, *Phys. Rev. Lett.* **84**, 5279 (2000).
3. G. Isidori, C. Smith, R. Underdorfer, *Eur. Phys. J. C* **36**, 57 (2004).
4. A. Alavi-Harati *for the KTeV Collaboration*, APS Division of Particles and Fields, Jan 5-9, 1999, UCLA, Los Angeles USA.
5. T.Andre, [hep-ph/0406006](#) and 6th International Conference on Hyperons, Charm and Beauty Hadrons, 27 June - 3 July 2004, Chicago USA.
6. A. Alavi-Harati *et al*, *Phys. Rev. D* **70**, 092007 (2004).
7. A. Alavi-Harati *et al*, *Phys. Rev. D* **71**, 012001 (2005).
8. A. Alavi-Harati *et al*, *Phys. Rev. D* **70**, 092006 (2004).
9. A. Alavi-Harati *et al*, *Phys. Rev. Lett.* **93**, 181802 (2004).
10. Particle Data Group, *Phys. Rev. D* **66**, 1 (2002).
11. V. Cirigliano, H. Neufeld, and H. Pichl, *Eur. Phys. J. C* **35**, 53 (2004).